

Synthesis of Type 1 Lewis b hexasaccharide antigen structures featuring flexible incorporation of L-[U-13C6]-fucose for NMR binding studies

Long, Mark; Ni Cheallaigh, Aisling; Reihill, Mark; Oscarson, Stefan; Lahmann, Martina

Organic and Biomolecular Chemistry

DOI:

[10.1039/D0OB00426J](https://doi.org/10.1039/D0OB00426J)

Published: 21/06/2020

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Long, M., Ni Cheallaigh, A., Reihill, M., Oscarson, S., & Lahmann, M. (2020). Synthesis of Type 1 Lewis b hexasaccharide antigen structures featuring flexible incorporation of L-[U-13C6]-fucose for NMR binding studies. *Organic and Biomolecular Chemistry*, 18(23), 4452-4458. <https://doi.org/10.1039/D0OB00426J>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?


Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Organic & Biomolecular Chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: M. B. Long, A. Ni Cheallaigh, M. Reihill, S. Oscarson and M. Lahmann, *Org. Biomol. Chem.*, 2020, DOI: 10.1039/D0OB00426J.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Synthesis of Type 1 Lewis b hexasaccharide antigen structures featuring flexible incorporation of L-[U-¹³C₆]-fucose for NMR binding studies

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Mark Long^a, Aisling Ni Cheallaigh^b, Mark Reihill^b, Stefan Oscarson^b, and Martina Lahmann^{a*}

While ¹³C-labelled proteins are common tools in NMR studies, lack of access to ¹³C-labelled carbohydrate structures has restricted their use. L-fucose is involved in a wide range of physiological and pathophysiological processes in mammalian organisms. Here, L-[U-¹³C₆]-Fuc labelled type I Lewis b (Le^b) structures have been synthesised for use in NMR binding studies with the Blood-group Antigen Binding Adhesin (BabA), a membrane-bound protein from the bacterium *Helicobacter pylori*. As part of this work, an efficient synthesis of a benzylated L-[U-¹³C₆]-Fuc thioglycoside donor from L-[U-¹³C₆]-Gal has been developed. The design and synthesis of an orthogonally protected tetrasaccharide precursor enabled controlled introduction of one or two ¹³C-labelled or non-labelled fucosyl residues prior to global deprotection. NMR analysis showed that it is straightforward to assign the anomeric centres as well as the H-5 positions to the individual fucosyl residues which are relevant for NMR binding studies.

Introduction

Helicobacter pylori is a spiral-shaped Gram-negative bacterium, which causes gastric and duodenal ulcers and is a major cause of stomach cancer. The bacterium selectively colonises the gastric epithelium.^{1–5} Unlike other pathogens, *Helicobacter pylori* has evolved to survive the highly acidic environment by metabolising urea to ammonia and carbon dioxide by a nickel depending urease.^{6–8} The bacterium expresses at least five types of adhesins⁹ which enable adherence to the stomach epithelium and concomitant production of several cytotoxins that destroy stomach epithelial cells, creating painful ulcers. The resulting chronic inflammation promotes cell proliferation, and thus predisposes the host to stomach cancer. One of the five adhesins expressed by *Helicobacter pylori* is the membrane-bound protein “Blood-group Antigen Binding Adhesin” (BabA). BabA binds to type I Lewis b (Le^b) antigens expressed on the surface of gastric epithelial cells. Analyses of binding specificities of *Helicobacter pylori* strains from across the world suggest that BabA has evolved in response to host mucosal

glycosylation patterns which permits *Helicobacter pylori* to adapt to its host and to maintain persistent colonisation.^{10–12}

A crystal structure of the carbohydrate-binding domain of BabA has been obtained while bound to a synthetic Lewis b structure. Analysis has revealed that the anchoring point for the binding of the glycan is the terminal disaccharide D-Galα(1→2) L-Fuc motif of the Le^b hexasaccharide.^{13, 14}

Detailed knowledge of these carbohydrate-protein interactions of *Helicobacter pylori* is crucial for understanding the structural and molecular basis for resulting diseases. While ¹³C-labelled proteins are common tools in NMR studies, lack of access to ¹³C-labelled carbohydrate structures has limited their use. Despite the difficulties arising from the need to synthesise the ¹³C-labelled material, ¹³C-enriched glycans are gaining traction as valuable tools for structure determination¹⁵ and conformational analysis¹⁶. Conformational analysis of a ligand bound to a lectin can be carried out by Saturation Transfer Difference (STD) and Transfer Nuclear Overhauser Effect (NOE) NMR experiments and do not require isotopically enriched material.¹⁷ Intramolecular NOE uses a combination of unlabelled carbohydrates and labelled proteins to extrapolate ligand-protein interactions^{18, 19}, while ¹³C-filtered NOESY can be used for identifying the conformation of bound ligands.^{20, 21} More recently, a combination of ¹³C-labelled carbohydrates and ¹⁵N (or ¹³C/¹⁵N) labelled proteins has been used to detail contact sites on the carbohydrate and the protein simultaneously.²² Availability of ¹³C-labelled Le^b structures would provide access to valuable tools for probing the interactions with BabA and other Le^b binding proteins.

Several strategies for the synthesis of type 1 Lewis structures have been reported. Danishefsky and co-workers have synthesised the Le^b tetrasaccharide and hexasaccharide using a polymer-based oligosaccharide preparation method with

^a M Long, M Lahmann
School of Natural Sciences
Bangor University
Bangor, Gwynedd, LL57 2UW, UK
E-mail: m.lahmann@bangor.ac.uk

^b A Ni Cheallaigh, M Reihill, S Oscarson
Centre for Synthesis and Chemical Biology
University College Dublin
Belfield, Dublin 4, IE

† Footnotes relating to the title and/or authors should appear here.


Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

ARTICLE

Journal Name

View Article Online
DOI: 10.1039/C9OB02613G

glycols as glycosyl donors.^{23, 24} Chernyak *et al.* have also reported the synthesis of the Le^b hexasaccharide using a convergent synthesis involving a tetrasaccharide thioglycoside donor and a spacer-equipped 3',4'-diol lactoside acceptor.²⁵ Furthermore, Lahmann *et al.* have reported the synthesis of the Le^b hexasaccharide, forming a tetrasaccharide acceptor *via* a [2+2] glycosylation which was then di-fucosylated.²⁶ Fournière *et al.* reported the synthesis of the Le^b pentasaccharide by first preparing the trisaccharide backbone, followed by di-fucosylation.²⁷ The Le^b tetrasaccharide was synthesised in addition to the Le^a trisaccharides by Ryzhov *et al.*²⁸ and Yan *et al.*²⁹ An enzymatic approach was also adopted by Chen and co-workers towards the truncated Le^b and Le^a structures.³⁰ Herein, we report the synthesis of a uniformly ¹³C labelled fucosyl donor as a general building block and an improved preparation of Lewis b structures *via* a linear approach. The flexibility of this approach allows access to novel mono and di L-[U-¹³C₆]-fucose labelled Le^b hexasaccharide structures, exemplified by the mono L-[U-¹³C₆]-Fuc-labelled Le^b hexasaccharide **1** and the di L-[U-¹³C₆]-Fuc-labelled Le^b hexasaccharide **2** (Scheme 1).




Scheme 1. The target structures and the building blocks shown in a simplified retrosynthesis scheme: mono L-[U-¹³C₆]-Fuc-labelled Le^b hexasaccharide **1**, di L-[U-¹³C₆]-Fuc-labelled Le^b hexasaccharide **2** and the unlabelled Le^b hexasaccharide **3** were prepared from the labelled and unlabelled building blocks **4** – **8**.

Results and discussion

The easiest access to isotopically enriched monosaccharides is via biosynthesis, which provides 100% uniformly [U-¹³C]

labelled material when the carbon source in the growth media is ¹³C labelled.³¹

In previous syntheses of Le^b hexasaccharide structures, a per-O-benzylated fucosyl thioglycoside donor has been used successfully, and therefore the analogous L-[U-¹³C₆]-Fuc **4** (Scheme 1) was envisioned as a suitable labelled donor. Since L-[U-¹³C₆]-Fuc is prohibitively expensive, the less costly L-[U-¹³C₆]-Gal was selected as starting material. The synthetic route was elaborated using unlabelled D-Gal (**S-24-S-29**, supplementary material) and then applied to L-[U-¹³C₆]-Gal (Scheme 2). Thus, after a near quantitative conversion of L-[U-¹³C₆]-Gal into its pentaacetate **9**^{32, 33}, a thiotolyl group was introduced at the anomeric position using *p*-thiocresol in the presence of BF₃·Et₂O (98% yield).³⁴ Deacetylation of **10**, followed by protection of the 6-position as TBDMS silyl ether and subsequent benzylation of the remaining free hydroxy groups, furnished L-galactoside **11** in a 75% yield over 3 steps. The silyl ether was removed with TBAF (**12**, 96% yield) and the 6-position was tosylated, producing a 94% yield of compound **13**. Reduction with LiAlH₄ in THF (61% yield) gave the desired L-[U-¹³C₆]-fucosyl donor **4** in an overall yield of 41% from L-[U-¹³C₆]-Gal.




Scheme 2. Reagents and conditions: a) 1. NaOAc, Ac₂O, 140 °C, 30 min, 99 % (**9**); 2. MePhSH, BF₃·Et₂O, CH₂Cl₂, rt, 2 h, 98% (**10**); b) 1. NaOMe, MeOH, rt, 30 min, 99%; 2. TBDMSCl, pyridine, 0 °C to rt, 4 h, 95%; 3. BnBr, NaH, DMF, 0 °C to rt, 2 h, 79% (**11**); c) Bu₄NF, THF, rt, 16 h, 96%; d) TsCl, pyridine, rt, 3 h, 94%; e) LiAlH₄, THF, reflux, 61%.

As expected, complex coupling patterns due to ¹³C-¹H and ¹³C-¹³C couplings were observed for the labelled L-Fuc donor **4** in the ¹H and ¹³C NMR spectra respectively but decoupling produced the matching spectra for the literature known unlabelled donor L-Fuc **5**³⁵ (Supplementary Figure 1).

The corresponding unlabelled L-Fuc **5**³⁵, the D-GlcNH₂ **7**³⁶, the D-Gal **8**³⁷ thioglycoside donors, and the D-Lac acceptor **6**³⁸ (Scheme 1) were prepared according to literature procedures. The assembly of the tetrasaccharide backbone **15** started with the NIS/AgOTf-mediated glycosylation at room temperature with D-Lac acceptor **6** and D-GlcNH₂ donor **7** to give the corresponding 1,2-*trans* linked trisaccharide **14**³⁸ (58%) (Scheme 3). The phthalimido and acetyl protecting groups were

removed simultaneously with EDA in EtOH. Subsequent selective *N*-acetylation was performed with Ac₂O in MeOH or toluene/MeOH (2:1, v/v), furnishing 3''-OH acceptor **15** (76%). In previous studies, we observed orthoester formation as a common side product during the glycosylation of acceptor **15** with the 2-*O*-acetyl analogue of donor **8**. 2-Benzoate protected glycosyl donors are less prone to orthoester formation.^{39,40} The 2-OBz thioglycoside donor **8** was successfully trialled and underwent glycosylation with acceptor **15** promoted by NIS and AgOTf (cat.) to furnish tetrasaccharide **16** in an 81% yield.




Scheme 3. Reagents and conditions: a) **7**, NIS, AgOTf (cat.), 4 Å molecular sieves, CH₂Cl₂, rt, 20 min, 58%; b) 1. EDA, EtOH, reflux, 16 h; 2. Ac₂O, MeOH/toluene (2:1, v/v), rt, 30 min; 76% over 2 steps; c) **8**, NIS, AgOTf (cat.), 4 Å molecular sieves, CH₂Cl₂, rt, 20 min – 2 h, 81%.

The orthogonal protecting group pattern of tetrasaccharide **16** permits the preparation of three different acceptors, and thus provides flexibility with respect to the position and number of L-[U-¹³C₆]-Fuc residues to be incorporated. One step deprotection exposes either the 4''-OH group when using reductive ring opening conditions (acceptor **17**) or the 2'''-OH position under Zemplén conditions (acceptor **18**), whilst sequential deprotection of both positions provides the corresponding diol acceptor **19**.

For the synthesis of the fully protected 2'''-mono L-[U-¹³C₆]-Fuc labelled Le^b hexasaccharide **22** (Scheme 3), tetrasaccharide **16** was subjected to reductive ring-opening conditions with NaBH₃CN and HCl/Et₂O in THF to reveal the 4''-OH, giving acceptor **17** (85%).^{41,42} Acceptor **17** underwent glycosylation with L-Fuc donor **5** using Lemieux's halide-assisted methodology to ensure α-selectivity.⁴³ This produced pentasaccharide **20** in a moderate yield of 45%. The removal of the 2'''-benzoyl group under Zemplén conditions required extended reaction times and a slightly elevated temperature in order to unmask the second fucosylation site (**21**, 93%). The L-[U-¹³C₆]-Fuc residue was installed using donor **4** and Lemieux's halide-assisted glycosylation conditions, and the 2'''-mono L-[U-¹³C₆]-Fuc labelled Le^b hexasaccharide **22** was obtained in a 79% yield. A similar reaction sequence with stepwise introduction of the fucosyl residues was also carried out with acceptor **18** to give the unlabelled 2'''-fucosylated pentasaccharide (73%).

Subsequent reductive ring-opening of the benzylidene acetal (66%) and a repeated fucosylation with unlabelled L-Fuc donor **4** provided the fully protected, unlabelled Le^b hexasaccharide **24**, demonstrating the feasibility of this approach to access also the 4''- mono L-[U-¹³C₆]-Fuc labelled Le^b hexasaccharide.



Scheme 4. Reagents and conditions: a) NaBH₃CN, 2 M HCl/Et₂O, 3 Å molecular sieves, THF, rt, 40 min, 85%; b) Br₂, Et₄NBr, 4 Å molecular sieves, CH₂Cl₂/DMF (9:1, v/v), rt, 16 h (donor **5**: **20**, 45%; donor **4**: **22**, 79%; **4**: **23**, 52%; **5**: **24**, 80%); c) 1 M NaOMe/MeOH, MeOH, rt to 40 °C (**21**, 26 h, 93%) and 1 M NaOMe/MeOH, MeOH/CH₂Cl₂ 7:1, rt to 40 °C (**18**, 48 h, 73%) and (**19**, 64 h, 88%).


The di-L-[U-¹³C₆]-Fuc labelled hexasaccharide **23** and unlabelled Le^b hexasaccharide **24** are also accessible from diol acceptor **19** obtained from tetrasaccharide **17** by debenzoylation (88%, Scheme 4). Interestingly, this transesterification was even slower than the corresponding reaction on the mono fucosylated pentasaccharide **21**. As above, the unlabelled L-Fuc donor **5** was converted into the corresponding bromide in the presence of acceptor diol **19**, with the glycosylation facilitated by Et₄NBr to give the protected unlabelled hexasaccharide **24** in an 80% yield (Scheme 4). The same halide-assisted glycosylation conditions were applied to the L-[U-¹³C₆]-Fuc donor **4** and diol acceptor **19** to produce the di-L-[U-¹³C₆]-Fuc labelled hexasaccharide **23** in a moderate yield (52%).

ARTICLE

Journal Name

View Article Online

DOI: 10.1039/D0OB00426J



Scheme 5. Reagents and conditions: a) H_2 , Pd/C, 0.1 M aq. HCl, 1,4-dioxane/ H_2O (3:1, v/v), rt, 2 d, (**1**, 87%; **2**, 30%; **3**, 96%).

The three hexasaccharides **22**, **23** and **24** were globally deprotected by Pd-catalysed hydrogenolysis to give targets **1**, **2** and **3** respectively. A variety of hydrogenolysis conditions were attempted. The presence of aqueous HCl was found to be crucial for efficient progression of the reaction in all cases, while a range of solvents were tolerated. The use of a mixture of heterogeneous catalysts (Pd/C and $\text{Pd}(\text{OH})_2/\text{C}$) was found to be beneficial in similar unpublished work but not for the compounds presented here. Applying optimised conditions, the deprotections proceeded very well for both the 2''-mono L-[U- $^{13}\text{C}_6$]-Fuc hexasaccharide **22** and the unlabelled hexasaccharide **24**, giving the mono-labelled (**1**, 87%) and unlabelled (**3**, 96%) Le^b hexasaccharides in excellent yields. After deprotection of **23**, a low yield of the di-L-[U- $^{13}\text{C}_6$]-Fuc labelled hexasaccharide **2** (30%) was recorded.

^1H NMR and ^1H - ^{13}C -HSQC was used to assign most of the signals and a CLIP-HSQC experiment confirmed the presence of two α -Fuc linkages and β -linkages for the remaining four glycosidic bonds of the known unlabelled Le^b hexasaccharide **3** (Figure 1 A and B). 1D TOCSY spectra were generated by selective excitation of each of the six H-1 signals individually (Figure 1 C). In combination with the NMR data obtained for the mono-labelled hexasaccharide **1** (vide infra), this enabled us, to assign undoubtedly the anomeric and the H-5 signals - which appeared in the same region occupied by the anomeric signals, to the individual 2''' and the 4'' fucosyl residues. There was no substantial difference in the chemical shift between the methyl groups of the two fucosyl residues which are otherwise distinct signals. The ability to distinguish between the individual fucosyl residues is of importance for NMR binding experiments with the labelled material.




Figure 1. – **A:** ^1H NMR spectrum of the unlabelled Le^b hexasaccharide **3**; **B:** CLIP-HSQC of **3**; **C:** 1D TOCSY experiments for **3**. The anomeric signals were selectively excited to generate spectra for each individual residue and then compared to the overall ^1H NMR spectrum (bottom).

The ^1H NMR spectrum of di- ^{13}C -labelled hexasaccharide **2** shows the expected additional ^{13}C - ^1H couplings corresponding to the labelled fucosyl residues (Figure 2A) which collapsed in the ^{13}C - ^1H decoupled experiments, reassembling the corresponding spectrum of the unlabelled hexasaccharide **3**. A standard ^{13}C NMR experiment displays only the ^{13}C enriched fucosyl residues, clearly showing the ^{13}C - ^{13}C couplings, and makes it a

convenient experiment to verify the incorporation of labelled material (Figure 2B).

View Article Online
DOI: 10.1039/D0OB00426J




Figure 2. – **A:** ^1H NMR spectrum of di- α -L-[U- $^{13}\text{C}_6$] fucosylated **2** with ^{13}C - ^1H couplings visible (bottom, red), ^{13}C - ^1H decoupled spectrum of **2** (middle, blue), and for comparison the ^1H NMR spectrum of the unlabelled hexasaccharide **3** (top, green); **B:** ^{13}C NMR spectrum of the di- α -L-[U- $^{13}\text{C}_6$] fucosylated hexasaccharide **2**. Only the signals from the uniformly labelled fucosyl residues are visible; **C:** Direct comparison of the ^{13}C NMR spectrum of mono 2'''- α -L-[U- $^{13}\text{C}_6$] fucosylated hexasaccharide **1** (bottom, red) with the ^{13}C NMR spectrum of the di labelled hexasaccharide **2** (top, blue) allows unambiguous identification of the anomeric carbons of the fucosyl residues. There is no significant difference in the C-6 shifts between the fucosyl residues; **D:** The coupled (bottom, red) and ^{13}C - ^1H decoupled (top, blue) ^1H NMR spectra of the 2'''- α -L-[U- $^{13}\text{C}_6$] fucosylated hexasaccharide **1**. The stacked view allows unambiguous assignment of the anomeric and H-5 signals for the individual fucosyl residues.

The ^1H NMR spectrum of the mono-labelled hexasaccharide **1** shows again the expected ^{13}C - ^1H couplings from the 2'''- α -L-[U- $^{13}\text{C}_6$] Fuc residue, while the ^{13}C - ^1H decoupled spectrum overlays with the unlabelled hexasaccharide **3**. The presence of the ^{13}C - ^1H couplings makes the identification of the anomeric proton and the H-5 for the 2'''- α -1 connected Fuc residue straightforward (Figure 2D). This information can also be obtained by comparing the ^{13}C NMR data of di-labelled compound **2** to the ^{13}C NMR data for the 2'''' mono labelled compound **1**, which shows the anomeric carbon of the α -L-[U-

$^{13}\text{C}_6$] Fuc-(1 \rightarrow 2)- β -D-Gal linkage to appear downfield to the α -L-[U- $^{13}\text{C}_6$] Fuc-(1 \rightarrow 4)- β -D-GlcNAc bond (Figure 2C).

Experimental

All experimental data are available in the supplementary file.

Conclusions

ARTICLE

Journal Name

View Article Online

DOI: 10.1039/C3OB26315E

In this work, an efficient synthesis of a ^{13}C -labelled L-fucosyl donor was developed from L-[U- $^{13}\text{C}_6$]-Gal. The synthesis of the L-[U- $^{13}\text{C}_6$]-fucose donor **4** was carried out on a 500 mg scale in a yield of 41% over 8 steps. This uniformly ^{13}C labelled fucosyl donor is a valuable building block for the preparation of labelled oligosaccharides for NMR binding studies. Thus, L-[U- $^{13}\text{C}_6$]-Fuc labelled type I Lewis b (Le^b) structures have been synthesised for use in NMR binding studies with BabA, a membrane-bound protein from the bacterium *Helicobacter pylori*. An orthogonally protected tetrasaccharide **16** allowed the regioselective introduction of one or two ^{13}C labelled fucose residues producing the protected ^{13}C labelled hexasaccharides **22** and **23**. Subsequent hydrogenolysis afforded the differently ^{13}C labelled hexasaccharides **1-3**. NMR analysis showed that it is straightforward to distinguish between the differently positioned fucosyl residues in the oligosaccharides. Making those and related structures a valuable tool for NMR protein-carbohydrate binding studies.

Acknowledgements

Financial support from the European Social Fund (KESS 80300, PhD scholarship for Mark Long) and Science Foundation Ireland Awards 13/IA/1959 and 16/RC/3889 is gratefully acknowledged (Aisling Ní Cheallaigh, Mark Reihill).

References

1. J. Parsonnet, G. D. Friedman, D. P. Vandersteen, Y. Chang, J. H. Vogelstein, N. Orentreich and R. K. Sibley, *N Engl J Med*, 1991, **325**, 1127-1131.
2. A. Dubois, *Emerg Infect Dis*, 1995, **1**, 79-85.
3. N. Uemura, S. Okamoto, S. Yamamoto, N. Matsumura, S. Yamaguchi, M. Yamakido, K. Taniyama, N. Sasaki and R. J. Schlemper, *N Engl J Med*, 2001, **345**, 784-789.
4. S. Suerbaum and P. Michetti, *N Engl J Med*, 2002, **347**, 1175-1186.
5. C. Y. Kao, B. S. Sheu and J. J. Wu, *Biomed J*, 2016, **39**, 14-23.
6. D. L. Weeks, S. Eskandari, D. R. Scott and G. Sachs, *Science*, 2000, **287**, 482-485.
7. E. F. Miller and R. J. Maier, *J. Bacteriol.*, 2014, **196**, 3074-3081.
8. M. Campanale, E. Nucera, V. Ojetto, V. Cesario, T. A. Di Rienzo, G. D'Angelo, S. Pecere, F. Barbaro, G. Gigante, T. De Pasquale, A. Rizzi, G. Cammarota, D. Schiavino, F. Franceschi and A. Gasbarrini, *Dig. Dis. Sci.*, 2014, **59**, 1851-1855.
9. D. J. Evans, Jr. and D. G. Evans, *Helicobacter*, 2000, **5**, 183-195.
10. T. Boren, P. Falk, K. A. Roth, G. Larson and S. Normark, *Science*, 1993, **262**, 1892-1895.
11. D. Ilver, A. Arnqvist, J. Ogren, I. M. Frick, D. Kersulyte, E. T. Inceci, D. E. Berg, A. Covacci, L. Engstrand and T. Boren, *Science*, 1998, **279**, 373-377.
12. M. Aspholm-Hurtig, G. Dailide, M. Lahmann, A. Kalia, D. Ilver, N. Roche, S. Vikstrom, R. Sjostrom, S. Linden, A. Backstrom, C. Lundberg, A. Arnqvist, J. Mahdavi, U. J. Nilsson, B. Velapattino, R. H. Gilman, M. Gerhard, T. Alarcon, M. Lopez-Brea, T. Nakazawa, J. G. Fox, P. Correa, M. G. Dominguez-Bello, G. I. Perez-Perez, M. J. Blaser, S. Normark, I. Carlstedt, S. Oscarson, S. Teneberg, D. E. Berg and T. Boren, *Science*, 2004, **305**, 519-522.
13. N. Hage, T. Howard, C. Phillips, C. Brassington, R. Overman, J. Debreczeni, P. Gellert, S. Stolnik, G. S. Winkler and F. H. Falcone, *Sci Adv*, 2015, **1**, e1500315-e1500315.
14. K. Moonens, P. Gideonsson, S. Subedi, J. Bugaytsova, E. Romão, M. Mendez, J. Nordén, M. Fallah, L. Rakhimova, A. Shevtsova, M. Lahmann, G. Castaldo, K. Brännström, F. Coppens, A. W. Lo, T. Ny, J. V. Solnick, G. Vandenbussche, S. Oscarson, L. Hammarström, A. Arnqvist, D. E. Berg, S. Muyldermans, T. Borén and H. Remaut, *Cell Host Microbe*, 2016, **19**, 55-66.
15. C. Fontana, H. Kovacs and G. Widmalm, *J. Biomol. NMR*, 2014, **59**, 95-110.
16. M. D. Battistel, M. Shangold, L. Trinh, J. Shiloach and D. I. Freedberg, *J. Am. Chem. Soc.*, 2012, **134**, 10717-10720.
17. B. Meyer and T. Peters, *Angew. Chem. Int. Ed. Engl.*, 2003, **42**, 864-890.
18. C. A. Bewley, *Structure*, 2001, **9**, 931-940.
19. M. Schubert, S. Bleuler-Martinez, A. Butschli, M. A. Walti, P. Egloff, K. Stutz, S. Yan, M. Collot, J. M. Mallet, I. B. Wilson, M. O. Hengartner, M. Aepli, F. H. Allain and M. Kunzler, *PLoS Pathog*, 2012, **8**, e1002706.
20. A. Canales, J. Angulo, R. Ojeda, M. Bruix, R. Fayos, R. Lozano, G. Gimenez-Gallego, M. Martin-Lomas, P. M. Nieto and J. Jimenez-Barbero, *J. Am. Chem. Soc.*, 2005, **127**, 5778-5779.
21. L. Nieto, A. Canales, G. Gimenez-Gallego, P. M. Nieto and J. Jimenez-Barbero, *Chemistry*, 2011, **17**, 11204-11209.
22. G. Nestor, T. Anderson, S. Oscarson and A. M. Gronenborn, *J. Am. Chem. Soc.*, 2017, **139**, 6210-6216.
23. V. Behar and S. J. Danishefsky, *Angew. Chem. Int. Ed.*, 1994, **33**, 1468-1470.
24. S. J. Danishefsky, V. Behar, J. T. Randolph and K. O. Lloyd, *J. Am. Chem. Soc.*, 1995, **117**, 5701-5711.
25. A. Chernyak, S. Oscarson and D. Turek, *Carbohydr. Res.*, 2000, **329**, 309-316.
26. M. Lahmann, L. Bülow, P. Teodorovic, H. Gyback and S. Oscarson, *Glycoconjugate J.*, 2004, **21**, 251-256.
27. V. Fournière, L. Skantz, F. Sajtos, S. Oscarson and M. Lahmann, *Tetrahedron*, 2010, **66**, 7850-7855.
28. I. M. Ryzhov, E. Y. Korchagina, A. B. Tuzikov, I. S. Popova, T. V. Tyrtsh, G. V. Pazynina, S. M. Henry and N. V. Bovin, *Carbohydr. Res.*, 2016, **435**, 83-96.
29. L. Yan and D. Kahne, *J. Am. Chem. Soc.*, 1996, **118**, 9239-9248.
30. H. Yu, Y. Li, Z. Wu, L. Li, J. Zeng, C. Zhao, Y. Wu, N. Tasnima, J. Wang, H. Liu, M. R. Gadi, W. Guan, P. G. Wang and X. Chen, *Chem. Commun.*, 2017, **53**, 11012-11015.
31. F. V. Filipp, N. Sinha, L. Jairam, J. Bradley and S. J. Opella, *J. Magn. Reson.*, 2009, **201**, 121-130.
32. M. L. Wolfrom and H. B. Wood, *J. Am. Chem. Soc.*, 1949, **71**, 3175-3176.
33. M. Lahmann and J. Thiem, *Carbohydr. Res.*, 1997, **299**, 23-31.
34. R. J. Ferrier and R. H. Furneaux, *Carbohydr. Res.*, 1976, **52**, 63-68.
35. Z. Zhang, I. R. Ollmann, X.-S. Ye, R. Wischnat, T. Baasov and C.-H. Wong, *J. Am. Chem. Soc.*, 1999, **121**, 734-753.
36. Q. Wang, J. Xue and Z. Guo, *Chem. Commun.*, 2009, 5536-5537.

ARTICLE

View Article Online

DOI: 10.1039/D0OB00426J

Journal Name

37. Z. Wang, L. Zhou, K. El-Boubbou, X.-s. Ye and X. Huang, *J. Org. Chem.*, 2007, **72**, 6409-6420.
38. A. Sundgren, M. Lahmann and S. Oscarson, *Bellstein Journal of Organic Chemistry*, 2010, **6**, 704-708.
39. P. J. Garegg, P. Konradsson, I. Kvarnstrom, T. Norberg, S. C. T. Svensson and B. Wigilius, *Acta Chemica Scandinavica, Series B: Organic Chemistry and Biochemistry*, 1985, **B39**, 569-577.
40. P. J. Garegg and T. Norberg, *Acta Chemica Scandinavica, Series B: Organic Chemistry and Biochemistry*, 1979, **B33**, 116-118.
41. P. J. Garegg and H. Hultberg, *Carbohydr. Res.*, 1981, **93**, C10-C11.
42. P. J. Garegg and H. Hultberg, *Carbohydr. Res.*, 1982, **110**, 261-266.
43. R. U. Lemieux and H. Driguez, *J. Am. Chem. Soc.*, 1975, **97**, 4069-4075.